

Is Second Trimester Abortion pill Affordable?



A medication abortion pill is not just a single pill, but it is a combination of two medications prescribed by the physician those work together to abort an unwanted pregnancy and shed the uterine lining. It is a popular name for using two types of medicine such as Mifepristone and Misoprostol to terminate a pregnancy. Mifepristone is taken first to block the progesterone hormone which is necessary to bear pregnancy and causing the embryo to detach from the uterine wall. Misoprostol is taken within a day or two days after taking first one. It helps to cause cramping that empties the uterus same like a period. According to medical research, it is proved that these pills are 96% effective in the 8-9 weeks and 93% effective in the 9-10 weeks. **Abortion pill online** are effective and provide the desired results.

No doubt, abortion pills are highly effective and safe. There are rare cases, when these pills get fail. According to the Center for Disease Control and Prevention's abortion surveillance data, only one woman dies after 100,000 legal abortion performed by the professional. The minor risk of dying from an abortion when woman took these pills after 10 weeks of pregnancy otherwise these pills are safe for every woman. Only 4% women are hospitalized after taking medication abortion due to some serious complications. The incredibly small risk of dying from an abortion rises the longer someone is pregnant, and since medication abortions are only advisable up to 10 weeks along, that makes them even safer. These pills have some side effects such as Nausea, cramping, infection,etc. Your doctor or nurse may offer you some treatment like Zofran and Phenergan to deal with Nausea. You can also use heating pads at home to deal with the bleeding. You can consult Second Trimester Abortion Clinic New York to get better treatment.



Second Trimester Abortion Clinic New York

Have a Question? Someone is available to answer your sexual health related question by e-mail 24 hours 7 days a week at 24hourabortion@gmail.com



Orlando Women's Center

Please call (443) 300-7455

You can take painkiller like ibuprofen about 30 minutes before you take the misoprostol to get rid of cramps. You can also take anti-nausea medicine after discussion with the doctor.Don't take aspirin, because it can make you bleed more.

You can confirm about your pregnancy with the help of an ultrasound. If you are taking abortion pills to terminate a pregnancy and you do not prefer an ultrasound first, then there are chances of the undetected ectopic pregnancy. If you do not bleed and do not pass the tissue after taking the Misoprostol, then you might have an ectopic pregnancy. Further, if you have mild or severe back and belly pain, you feel you might faint or if you do so and have severe shoulder pain then you might have an ectopic pregnancy that has burst and you should consult doctor immediately. This type of pregnancy is treated everywhere, even at those places also where abortion is severely restricted.

Some women find that it helps to have a partner or friend with them when they use the misoprostol. Some women may prefer to have privacy, but have someone they know they can reach if they need help. While it is rare to have an emergency, it is important that you make plans ahead of time before seeking medical attention. F you are more than 10 weeks pregnant follow Second trimester abortion procedures.

You cannot get the pills over the counter as these are considered as a prescription drug and can only be provided to the consumer by the doctor prescribed. Furthermore, when come to

know how to get these abortion pills, you should be aware that these pills cannot be taken legally unless your doctor suggest you for this. Abortion pills are the fastest and easiest way to get rid of unexpected or unwanted pregnancy. However, abortion not only drains a woman physically it exhausts her mentally too. Moreover, Second trimester abortion pill is easy to afford and every person can easily afford it.